

RESOURCES

Philadelphia Department of Public Health

overdose.prevention@phila.gov
www.PhillyNaloxone.com

DBHIDS Warm Line

Talk with trained, certified, supportive peer specialists.
Monday–Friday 4PM-7PM
(855) 507-WARM (9276)

Get more fentanyl test strips, naloxone (Narcan), or harm reduction info

- Prevention Point
- www.PhillyNaloxone.com
- www.NextDistro.org/Philly

MOBILE SITES

Prevention Point

Call for times and locations.
2913-15 Kensington Ave
Philadelphia, PA 19134
(215) 530-2370
ppponline.org

Services available

- OD prevention education
- Naloxone (Narcan)
- Syringe Exchange
- HCV and HIV screening
- Medical Triage
- Fentanyl test strips

USING ALONE?

Remote Supervision Resources

Anonymously connect with a peer who can contact a friend for you or call 911 in case you become unresponsive:

Never Use Alone Hotline:

E: 1-800-484-3731

S: 1-800-928-5330

The Brave App:

www.thebraveapp.com

In Apple and Google Play store



**KNOW
WHAT'S IN
YOUR DRUGS**

HOW TO USE FENTANYL TEST STRIPS

Test Strip Directions

- 1) Prepare your drugs. You only need a small amount for the test. Residue from bag/cooker works.
- 2) Add fresh water to empty bag/cooker you just prepped. Mix well.
- 3) Dip wavy-line end of test strip in water for 15 seconds (don't pass solid blue line).
- 4) Remove and lay on flat surface. Results are usually ready within 1 minute.

Testing Pills

If you take pills orally:

- Scrape a small amount off the pill or cut pill in half
- Scrape from middle for best results.
- Follow test strip directions.

If you snort:

- Crush pills and add water.
- You can also use a few grains from pill, but testing residue is safer.

Test Results





One line = **Positive** for fentanyl.



Two lines = **Negative** for fentanyl.

If second line (closest to white end) is faint, it is negative for fentanyl.

PLEASE NOTE:

- Test is invalid if:
 - 1) test has no lines OR
 - 2) red line closest to blue end is not there.
- Try a small amount first and go slow when using.

Fentanyl may still be present even if the test is negative.



Always carry naloxone!

